

Spatial assignment of test sample

December 14, 2016

Contents

Input	1
Isotope values of test sample	1
Output	1
Model	1
Map of best fitted reference sample	2
Best fitted reference entries	3
Testing robustness of assignment: Wilcoxon signed rank test	3
P-values for the k nearest neighbours in Wilcoxon Test	3
Goodness of fit of test sample:	4

Input

Website Identifier: 2

Isotope values of test sample

Table 1: Isotope values of test sample


13C/12C	15N/14N	18O/16O	2H/1H	34S/32S
-20.5	7.6	16	-30.8	8.2

Output

Model

```
##  
## Call:  
## train.kknn(formula = fmla, data = ivory.train, kmax = 15, distance = 2, kernel = knl)  
##  
## Type of response variable: nominal  
## Minimal misclassification: 0.3765867  
## Best kernel: triangular  
## Best k: 15  
  
Classifier: country_code
```


Map of best fitted reference sample



Best fitted reference sample:

- Site: Dem. Rep. Congo, Gangala
- Country: CD
- Region: Central Africa
- CITES: Appendix I
- Lat: 3.71
- Lon: 29.12

Assignment of test sample to nearest neighbours



Best fitted reference entries

Table 2: Details of best fitted reference entry (row 1) and other fitted entries within best classifier

lon	lat	location	13C/12C	15N/14N	18O/16O	2H/1H	34S/32S
29.12	3.71	Dem. Rep. Congo, Gangala	-20.1	7.5	17.0	-33.0	7.1
29.09	3.40	Dem. Rep. Congo, Gangala na Bodio	-19.7	8.2	17.5	-29.9	7.7
23.05	-7.19	Dem. Rep. Congo, Lula riv	-20.9	6.9	17.8	-30.6	7.7
29.09	3.40	Dem. Rep. Congo, Gangala na Bodio	-20.6	8.2	16.7	-29.0	5.0
23.00	4.00	Dem. Rep. Congo, Bas Uele	-20.8	7.9	18.0	-35.6	8.8
27.20	-2.40	Dem. Rep. Congo, Shabunda terr	-20.0	9.5	16.9	-27.7	8.4

Country of prediction: CD

Testing robustness of assignment: Wilcoxon signed rank test

If p-value > 0.05, the test concludes that the isotope signature of the test sample is similar to the respective nearest neighbour reference sample.

P-values for the k nearest neighbours in Wilcoxon Test

“0.65162, 0.65162, 0.01784, 0.01048, 0.00191, 0.00028”

Goodness of fit of test sample:

- good fit: if $p > 0.05$ for at least two tested nearest neighbour reference samples;
- moderate fit: if $p > 0.05$ for at least one tested nearest neighbour reference samples;
- uncertain: if $p > 0.05$ for none of the tested nearest neighbour reference samples.

Assumption: At least two nearest reference samples are available.

Overall goodness of fit of test sample: “**good fit**”