

# Spatial assignment of test sample

November 24, 2016

## Contents

<b>Input</b>	<b>1</b>
Isotope values of test sample . . . . .	1
<b>Output</b>	<b>1</b>
Model . . . . .	1
Map of best fitted reference sample . . . . .	2
Best fitted reference entries . . . . .	3
Testing robustness of assignment: Wilcoxon signed rank test . . . . .	3
P-values for the k nearest neighbours in Wilcoxon Test . . . . .	3
Goodness of fit of test sample: . . . . .	3

## Input

Website Identifier:

### Isotope values of test sample

Table 1: Isotope values of test sample


13C/12C	15N/14N	18O/16O	2H/1H	34S/32S
-22.7	10.6	15.9	-59.2	8.6

## Output

### Model

```
##
## Call:
## train.kknn(formula = fmla, data = ivory.train, kmax = 15, distance = 2, kernel = knl)
##
## Type of response variable: nominal
## Minimal misclassification: 0.3765867
## Best kernel: triangular
## Best k: 15
Classifier: country_code
```


## Map of best fitted reference sample



### Best fitted reference sample:

- Site: Tanzania, Gebirgswald im Ngorongoro
- Country: TZ
- Region: East Africa
- CITES: Appendix I
- Lat: -3.205556
- Lon: 35.463611

## Assignment of test sample to nearest neighbours



### Best fitted reference entries

Table 2: Details of best fitted reference entry (row 1) and other fitted entries within best classifier

lon	lat	location	13C/12C	15N/14N	18O/16O	2H/1H	34S/32S
35.46	-3.21	Tanzania, Gebirgswald im Ngorongoro	-22.7	9.1	15.6	-61.3	9.2
35.46	-3.21	Tanzania, Gebirgswald im Ngorongoro	-21.3	10.1	15.0	-59.3	9.1
35.46	-3.00	Tanzania, Gebirgswald im Ngorongoro	-21.0	10.1	15.4	-61.0	6.8
35.33	-7.20	Tanzania	-22.1	9.5	16.5	-56.7	5.4

Country of prediction: **TZ**

### Testing robustness of assignment: Wilcoxon signed rank test

If  $p\text{-value} > 0.05$ , the test concludes that the isotope signature of the test sample is similar to the respective nearest neighbour reference sample.

### P-values for the k nearest neighbours in Wilcoxon Test

“0.331, 0.310, 0.172, 0.037”

### Goodness of fit of test sample:

- good fit: if  $p > 0.05$  for at least two tested nearest neighbour reference samples;

- moderate fit: if  $p > 0.05$  for at least one tested nearest neighbour reference samples;
- uncertain: if  $p > 0.05$  for none of the tested nearest neighbour reference samples.

Assumption: At least two nearest reference samples are available.

Overall goodness of fit of test sample: “**good fit**”