

Spatial assignment of test sample

December 13, 2016

Contents

Input	1
Isotope values of test sample	1
Output	1
Model	1
Map of best fitted reference sample	2
Best fitted reference entries	3
Testing robustness of assignment: Wilcoxon signed rank test	3
P-values for the k nearest neighbours in Wilcoxon Test	3
Goodness of fit of test sample:	4

Input

Website Identifier: 88

Isotope values of test sample

Table 1: Isotope values of test sample


13C/12C	15N/14N	18O/16O	2H/1H	34S/32S
-21	7	16.6	-44.3	8

Output

Model

```
##  
## Call:  
## train.kknn(formula = fmla, data = ivory.train, kmax = 15, distance = 2, kernel = knl)  
##  
## Type of response variable: nominal  
## Minimal misclassification: 0.3765867  
## Best kernel: triangular  
## Best k: 15  
  
Classifier: country_code
```


Map of best fitted reference sample



Best fitted reference sample:

- Site: Mozambique, Kambako
- Country: MZ
- Region: Southern Africa
- CITES: Appendix I
- Lat: -14.75
- Lon: 32

Assignment of test sample to nearest neighbours



Best fitted reference entries

Table 2: Details of best fitted reference entry (row 1) and other fitted entries within best classifier

lon	lat	location	13C/12C	15N/14N	18O/16O	2H/1H	34S/32S
32.00	-14.75	Mozambique, Kambako	-20.8	7.8	16.0	-45.0	9.8
39.00	-11.18	Mozambique, Rovuma river area	-20.8	7.0	15.4	-40.0	7.5
31.15	-14.86	Mozambique, Bairro Gebeuza village	-19.7	7.3	16.4	-46.3	8.9
37.84	-12.31	Mozambique, Block L7 - Lugenda south ban	-22.1	8.0	16.2	-44.5	7.6
37.84	-12.30	Mozambique, Block L7 - Lugenda south ban	-22.0	7.5	15.5	-41.6	8.4
37.30	-12.00	Mozambique, Luwure (block L7) area	-22.5	6.7	16.5	-41.6	8.9

Country of prediction: MZ

Testing robustness of assignment: Wilcoxon signed rank test

If p-value > 0.05, the test concludes that the isotope signature of the test sample is similar to the respective nearest neighbour reference sample.

P-values for the k nearest neighbours in Wilcoxon Test

“0.13370, 0.10231, 0.08476, 0.05677, 0.05677, 0.00014”

Goodness of fit of test sample:

- good fit: if $p > 0.05$ for at least two tested nearest neighbour reference samples;
- moderate fit: if $p > 0.05$ for at least one tested nearest neighbour reference samples;
- uncertain: if $p > 0.05$ for none of the tested nearest neighbour reference samples.

Assumption: At least two nearest reference samples are available.

Overall goodness of fit of test sample: “**good fit**”