Spatial assignment of test sample

November 24, 2016

Contents

| Inp | ut | 1 |
|-----|---|---|
| Ī | Isotope values of test sample | 1 |
| Out | tput | 1 |
| I | Model | 1 |
| I | Map of best fitted reference sample | 2 |
|] | Best fitted reference entries | 3 |
| - | Testing robustness of assignment: Wilcoxon signed rank test | 3 |
| | P-values for the k nearest neighbours in Wilcoxon Test | |
| | Goodness of fit of test sample: | 4 |

Input

Website Identifier: $T2_39cm$

Isotope values of test sample


Table 1: Isotope values of test sample

| 13C/12C | 15N/14N | 18O/16O | $2\mathrm{H}/1\mathrm{H}$ | 34S/32S |
|---------|---------|---------|---------------------------|---------|
| -19.8 | 6.2 | 19 | -48.1 | 8.9 |

Output

Model

Map of best fitted reference sample




Best fitted reference sample:

• Site: Southern Zambia

• Country: ZM

Region: Southern Africa
CITES: Appendix I
Lat: -14.972185
Lon: 25.957587

Assignment of test sample to nearest neighbours



Best fitted reference entries

Table 2: Details of best fitted reference entry (row 1) and other fitted entries within best classifier

| lon | lat | location | 13C/12C | $15\mathrm{N}/14\mathrm{N}$ | 18O/16O | 2H/1H | 34S/32S |
|-------|--------|--|---------|-----------------------------|---------|-------|---------|
| 25.96 | -14.97 | Southern Zambia | -20.4 | 7.0 | 20.1 | -49.2 | 8.5 |
| 25.96 | -14.97 | Southern Zambia | -20.6 | 7.1 | 19.6 | -51.9 | 8.2 |
| 32.10 | -12.05 | Southern Zambia | -19.6 | 7.3 | 18.3 | -45.3 | 7.4 |
| 32.19 | -11.41 | Southern Zambia | -20.6 | 7.2 | 18.4 | -46.4 | 7.3 |
| 25.45 | -16.52 | Southern Zambia | -21.3 | 7.1 | 18.8 | -49.7 | 8.9 |
| 31.44 | -12.12 | Northeastern Zambia, near Chilonga | -20.9 | 6.9 | 18.7 | -53.0 | 7.4 |
| 32.38 | -11.29 | North Zambia, near Msitu | -20.7 | 7.0 | 18.2 | -44.8 | 10.6 |
| 32.30 | -11.37 | Southern Zambia | -20.8 | 7.3 | 18.6 | -54.2 | 9.6 |
| 31.21 | -13.62 | Zambia, Luangwa River, western of Chipat | -18.7 | 7.5 | 19.3 | -43.7 | 9.9 |

Country of prediction: ZM

Testing robustness of assignment: Wilcoxon signed rank test

If p-value > 0.05, the test concludes that the isotope signature of the test sample is similar to the respective nearest neighbour reference sample.

P-values for the k nearest neighbours in Wilcoxon Test

 $\begin{tabular}{l} "0.376588383, \ 0.084761154, \ 0.002675546, \ 0.001114922, \ 0.000515506, \ 0.000111219, \ 0.000003584, \ 0.000002501, \ 0.000000026" \end{tabular}$

Goodness of fit of test sample:

- good fit: if p > 0.05 for at least two tested nearest neighbour reference samples;
- moderate fit: if p > 0.05 for at least one tested nearest neighbour reference samples;
- uncertain: if p > 0.05 for none of the tested nearest neighbour reference samples.

Assumption: At least two nearest reference samples are available.

Overall goodness of fit of test sample: "good fit"