Spatial assignment of test sample

November 24, 2016

Contents

Inpu	ıt	1
Is	sotope values of test sample	1
Out	put	1
N	Iodel	1
N	Iap of best fitted reference sample	2
В	Sest fitted reference entries	3
Τ	esting robustness of assignment: Wilcoxon signed rank test	3
	P-values for the k nearest neighbours in Wilcoxon Test	
	Goodness of fit of test sample:	4

Input

Website Identifier: $T2_33cm$

Isotope values of test sample

Table 1: Isotope values of test sample


13C/12C	$15\mathrm{N}/14\mathrm{N}$	18O/16O	$2\mathrm{H}/1\mathrm{H}$	34S/32S
-19.8	6	18.5	-51.1	9

Output

Model

```
##
## Call:
## train.kknn(formula = fmla, data = ivory.train, kmax = 15, distance = 2, kernel = knl)
##
## Type of response variable: nominal
## Minimal misclassification: 0.3765867
## Best kernel: triangular
## Best k: 15
Classifier: country_code
```

Map of best fitted reference sample




Best fitted reference sample:

• Site: Southern Zambia

• Country: ZM

Region: Southern Africa
CITES: Appendix I
Lat: -16.093781
Lon: 25.880879

Assignment of test sample to nearest neighbours



Best fitted reference entries

Table 2: Details of best fitted reference entry (row 1) and other fitted entries within best classifier

lon	lat	location	13C/12C	15N/14N	18O/16O	$2\mathrm{H}/1\mathrm{H}$	34S/32S
25.88	-16.09	Southern Zambia	-21.0	6.4	18.1	-56.6	9.6
25.68	-16.76	Southern Zambia	-20.7	6.4	18.0	-57.7	10.0
31.44	-12.12	Northeastern Zambia, near Chilonga	-20.9	6.9	18.7	-53.0	7.4
32.30	-11.37	Southern Zambia	-20.8	7.3	18.6	-54.2	9.6
25.96	-14.97	Southern Zambia	-20.6	7.1	19.6	-51.9	8.2
25.88	-16.09	Southern Zambia	-20.8	6.5	17.8	-56.3	10.5
25.45	-16.52	Southern Zambia	-21.3	7.1	18.8	-49.7	8.9
32.10	-12.05	Southern Zambia	-19.6	7.3	18.3	-45.3	7.4
32.19	-11.41	Southern Zambia	-20.6	7.2	18.4	-46.4	7.3
26.07	-15.90	Southern Zambia	-19.7	7.7	17.9	-54.9	9.9
25.96	-14.97	Southern Zambia	-20.4	7.0	20.1	-49.2	8.5

Country of prediction: ZM

Testing robustness of assignment: Wilcoxon signed rank test

If p-value > 0.05, the test concludes that the isotope signature of the test sample is similar to the respective nearest neighbour reference sample.

P-values for the k nearest neighbours in Wilcoxon Test

 $\begin{tabular}{l} "0.201204919, \ 0.002675546, \ 0.002262130, \ 0.000764607, \ 0.000515506, \ 0.000068155, \ 0.000017612, \ 0.000003584, \ 0.000000490, \ 0.000000490, \ 0.000000026" \end{tabular}$

Goodness of fit of test sample:

- good fit: if p > 0.05 for at least two tested nearest neighbour reference samples;
- moderate fit: if p > 0.05 for at least one tested nearest neighbour reference samples;
- uncertain: if p > 0.05 for none of the tested nearest neighbour reference samples.

Assumption: At least two nearest reference samples are available.

Overall goodness of fit of test sample: " $\mathbf{moderate}$ fit"